Amendmen After Decision on Appeal

Serial No. 09/027,439

BEST AVAILABLE COPY

This listing of claims will replace all prior versions, and listings, of claims in the application:

LISTING OF CLAIMS:

1.-4". (Canceled)

48. (Previously Presented) An isolated nucleic acid molecule consisting of SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, or SEQ ID NO: 6, or an RNA equivalent thereof,

or a nucleic acid complementary to said isolated molecule, capable of base-pairing according to the standard Watson-Crick complementarity rules,

or a nucleic acid substantially complementary to said isolated molecule which is capable of h/bridizing to the nucleic acid molecule under the following stringent conditions:

hybridization at 40°-65°C for 14-16 hours in a hybridization solution at ph 7.8, containing 0.9 M NaCl, 0.12 M Tris-HCl, 6nM EDTA, 0.1M sodium phosphate buffer, 0.1% SDS and 0.1% polyvinylpyrrolidone,

followed by three 15-minute washes at 40°-65°C to remove unbound probes in a solution at pH 7, containing 0.075 M NaCl, 0.0075 M Na Citrate and 0.1% SDS.

49.-51. (Canceled)

- 52. (Previously Presented) The isolated nucleic acid molecule consisting of the nucleotice sequence of SEQ ID NO: 6.
- 53. (Previously Presented) An isolated nucleic acid molecule comprising a nucleotide sequence of SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, or SEQ ID NO: 6,

or an RNA equivalent thereof,

or a nucleic acid complementary to said isolated molecule, capable of base-pairing according to the standard Watson-Crick complementarity rules.

- 54. (Cancelled)
- 55. (Previously Presented) A probe which

Amendmen: After Decision on Appeal

Serial No. 09/027,439

BEST AVAILABLE COPY

- a) targets Shigella flexneri comprising a fragment greater than 10 to 40 bases in length of a nucleotide sequence SEQ ID NO: 3, an RNA equivalent thereof, or a nucleic acid complementary to said molecule, capable of base-pairing according to the standard Watson-Crick complementarity rules,
- b) targets Shigella sonnei comprising a fragment greater than 10 to 40 bases in length of a nucleotide sequence SEQ ID NO: 4, an RNA equivalent thereof, or a nucleic acid complementary to said molecule, capable of base-pairing according to the standard Watson-Crick complementarity rules,
- c) targets Shigella dysenteriae comprising a fragment greater than 10 to 40 bases in length of a nucleotide sequence SEQ ID NO: 5, an RNA equivalent thereof, or a nucleic acid complementary to said molecule, capable of base-pairing according to the standard Watson-Crick complementarity rules,

or

d) targets Shigella boydii comprising a fragment greater than 10 to 40 bases in length of a rucleotide sequence SEQ ID NO: 6, an RNA equivalent thereof, or a nucleic acid complementary to said molecule, capable of base-pairing according to the standard Watson-Crick complementarity rules.

56. (Previously Presented) A probe which

- a) targets Shigella flexners consisting of a fragment greater than 10 to 40 bases in length of a rucleotide sequence SEQ ID NO: 3, an RNA equivalent thereof, or a nucleic acid complementary to said molecule, capable of base-pairing according to the standard Watson-Crick complementarity rules,
- b) targets Shigella sonnei consisting of a fragment greater than 10 to 40 bases in length of a rucleotide sequence SEQ ID NO: 4, an RNA equivalent thereof, or a nucleic acid complementary to said molecule, capable of base-pairing according to the standard Watson-Crick complementarity rules,
- c) targets Shigella dysenteriae consisting of a fragment greater than 10 to 40 bases in length of a nucleotide sequence SEQ ID NO: 5, an RNA equivalent thereof, or a nucleic acid complementary to said molecule, capable of base-pairing according to the standard Watson-Crick complementarity rules,

Or

Amendment After Decision on Appeal

Serial No. 09/027,439

BEST AVAILABLE COPY

- d) targets Shigella boydit consisting of a fragment greater than 10 to 40 bases in length of a nucleotide sequence SEQ ID NO: 6, an RNA equivalent thereof, or a nucleic acid complementary to said molecule, capable of base-pairing according to the standard Watson-Crick complementarity rules.
- 57. (Previously Presented) A probe as in claim 55 which comprises 15-25 bases in length.
- 58. (Previously Presented) A probe as in claim 56 which comprises 15-25 bases in length.